



Physiological characterization of three different *Yarrowia lipolytica* strains growing on glycerol, glucose, xylose and arabinose

Lubuta, Patrice; Workman, Mhairi

Published in:
Book of Abstracts. DTU's Sustain Conference 2015

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Lubuta, P., & Workman, M. (2015). Physiological characterization of three different *Yarrowia lipolytica* strains growing on glycerol, glucose, xylose and arabinose. In *Book of Abstracts. DTU's Sustain Conference 2015* [B-12] Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Physiological characterization of three different *Yarrowia lipolytica* strains growing on glycerol, glucose, xylose and arabinose

Patrice Lubuta, Mhairi Workman*

Section for Eukaryotic Biotechnology, DTU Systems Biology

*Corresponding author email: mwo@bio.dtu.dk

The nonconventional yeast *Yarrowia lipolytica* is, due to favorable physiological features, a promising candidate for future biotechnological applications. As an oleaginous yeast *Y. lipolytica* is able to synthesize and accumulate lipids to levels exceeding 50% of cell dry weight (Beopoulos et al., 2009). Besides that, this species can secrete large amounts of organic acids (e.g. citric and isocitric acids, α -ketoglutaric acid), produce sugar alcohols (e.g. mannitol, erythritol) and single-cell protein. It can also be used as a platform for the production of various heterologous proteins (e.g. lipases, proteases) (Darvishi Harzevili, 2014). Furthermore *Y. lipolytica* is generally recognized as safe (GRAS). Besides the industrial applications, *Y. lipolytica* is also used as a model organism in several studies of academic interest (Barth, 2013).

The overall goal of this research project is to evaluate *Y. lipolytica*'s potential to be used as a biocatalyst in a biorefinery process where the conversion of agro-industrial wastes and renewable low-cost substrates (such as glycerol) to value added products is desired. A challenge in this context is the seasonal variation in availability and composition of biomass. This study addresses, therefore, the applicability of *Y. lipolytica* as a cell factory for conversion of glycerol and hydrolysates of lignocellulosic materials which mainly consists of fermentable sugars like xylose, glucose and arabinose (Tsigie et al., 2012). A quantitative physiological investigation was carried out in order to assess the cellular performance on different carbon sources and on combinations of these. In this context we were also interested in strain diversity regarding carbon utilization. The DTU in-house strain IBT 446, and the publically available strains W29 and H222 were used in this study.

Barth, G. (Ed.). (2013). *Yarrowia lipolytica* (Vol. 25). Berlin, Heidelberg: Springer Berlin Heidelberg. doi:10.1007/978-3-642-38583-4

Beopoulos, A., Cescut, J., Haddouche, R., Uribelarrea, J.-L., Molina-Jouve, C., & Nicaud, J.-M. (2009). *Yarrowia lipolytica* as a model for bio-oil production. *Progress in Lipid Research*, 48(6), 375–87. doi:10.1016/j.plipres.2009.08.005

Darvishi Harzevili, F. (2014). *Biotechnological Applications of the Yeast Yarrowia lipolytica*. Cham: Springer International Publishing. doi:10.1007/978-3-319-06437-6

Tsigie, Y. A., Wang, C.-Y., Kasim, N. S., Diem, Q.-D., Huynh, L.-H., Ho, Q.-P., ... Ju, Y.-H. (2012). Oil production from *Yarrowia lipolytica* Po1g using rice bran hydrolysate. *Journal of Biomedicine & Biotechnology*, 2012, 378384. doi:10.1155/2012/378384

